

Dissipative quantum dynamics of bosonic atoms in a shallow 1D optical lattice

J. Ruostekoski^{1,2} and L. Isella¹

¹*Department of Physics, Astronomy and Mathematics,
University of Hertfordshire, Hatfield, Herts, AL10 9AB, UK*

²*Institute for Theoretical Atomic and Molecular Physics,
Harvard-Smithsonian Center for Astrophysics, Cambridge MA 02135*

(Dated: February 2, 2008)

We theoretically study the dipolar motion of bosonic atoms in a very shallow, strongly confined 1D optical lattice using the parameters of the recent experiment [Fertig *et al.*, Phys. Rev. Lett. **94**, 220402 (2005)]. We find that, due to momentum uncertainty, a small, but non-negligible, atom population occupies the unstable velocity region of the corresponding classical dynamics, resulting in the observed dissipative atom transport. This population is generated even in a static vapor, due to quantum fluctuations which are enhanced by the lattice and the confinement, and is not notably affected by the motion of atoms or finite temperature.

PACS numbers: 03.75.Lm, 03.75.Kk, 03.75.Gg

The rich phenomenology of the dynamics of an atomic Bose-Einstein condensate (BEC) in a periodic optical lattice potential has recently attracted considerable theoretical and experimental interest. In particular, the non-linear mean-field interaction of the BEC may give rise to dynamical and energetic instabilities in the transport properties of the atoms [1, 2, 3, 4, 5, 6, 7, 8, 9, 10]. Most experiments to date have been carried out in shallow lattices where the motion of the BEC can be accurately described by the classical mean-field theories. In the strongly interacting regime, with very high lattice potentials, the atoms were shown to undergo a quantum phase transition to the Mott insulator state [11, 12]. Recently strongly damped oscillations of bosonic atoms were experimentally observed by Fertig *et al.* [13] at NIST in a tight elongated 1D atom trap, even in the limit of a very shallow lattice potential. In this Letter we numerically investigate the damped oscillations of bosonic atoms in a shallow lattice using the parameters of Ref. [13] and show that the observed damping rate can be explained by the growing population in dynamically unstable velocity region, resulting in phase decoherence. This population appears even without the dipolar motion and is not notably affected in the studied case by the atom dynamics. Moreover, for fixed nonlinearity, both, the population at unstable velocity and the damping rate, have approximately exponential dependence on the atom number. For larger atom numbers the quantum fluctuations are suppressed and the finite-temperature effects can become observable.

In the recent NIST experiment [13] an array of independent 1D atom ‘tubes’ were generated by applying a strong transverse 2D optical lattice potential. The bosonic atoms in each tube were also trapped along the axial direction by a harmonic magnetic potential and by a very shallow periodic optical lattice. The dipole oscillations of atoms along the weak axial lattice were excited by suddenly displacing the harmonic trap by means of

applying a linear magnetic field gradient. In the absence of the lattice, the dipolar motion is undamped and unaffected by the interactions. However, significantly inhibited dynamics was observed even in the presence of a very shallow lattice with the height of $0.25E_r$, where $E_r \equiv 2\pi^2\hbar^2/m\lambda^2$ is the lattice photon recoil energy with the wavelength λ . The strong effect of such a shallow lattice on dissipation is surprising, since the lattice modulates the atom density only by 6%. Due to the limited resolution of the imaging system, it was not possible to measure the damping of the oscillations in the center-of-mass (cm) position of the atoms, but the oscillations in the cm velocity were imaged after a time-of-flight, following the sudden turn-off of the trapping potentials.

Here we numerically study the dipole oscillations of bosonic atoms in a 1D lattice within the truncated Wigner approximation (TWA) for the complete multi-mode field operator, beyond the discrete tight-binding approximation. The previous theoretical studies of the quantum dynamics of bosonic atoms in a combined harmonic and lattice potential indicate that quantum fluctuations can result in strong dissipation [14, 15, 16, 17]. In Ref. [14] the dynamics was studied within the tight-binding approximation to the TWA for deep lattices. This approach is much less accurate for very shallow lattices studied here and in Ref. [13]. Moreover, we find that the tight-binding approximation to the TWA for shallow lattices strongly underestimates the generated noise and the effect of vacuum fluctuations on the dynamics.

In Refs. [13, 18] the atoms were confined in a 2D array of decoupled tight 1D tubes by means adiabatically loading the atoms into a strong 2D optical lattice. We write the atom density in a 1D tube at (x_0, y_0) :

$$\rho(\mathbf{r}) = \frac{15N_a d^2}{8\pi^2 l_\perp^2 \bar{R}^3} \left(1 - \frac{x_0^2}{\bar{R}_x^2} - \frac{y_0^2}{\bar{R}_y^2} - \frac{z^2}{\bar{R}_z^2}\right) e^{-\bar{\rho}^2/l_\perp^2}, \quad (1)$$

where $\bar{\rho}^2 \equiv (x-x_0)^2 + (y-y_0)^2$, $\bar{R}_i$ are the Thomas-Fermi radii of the 3D BEC, with $\bar{R} \equiv (\bar{R}_x \bar{R}_y \bar{R}_z)^{1/3}$, N_a denotes

the total atom number, and $d = \lambda/2$ the lattice spacing. By expanding around the lattice site minimum we may estimate $l_\perp \simeq (\hbar/m\omega_\perp)^{1/2}$ by the harmonic trap frequency in the transverse direction $\omega_\perp = 2\sqrt{s}E_r/\hbar$, where s denotes the lattice height in the units of E_r . In Ref. [13], $s = 30$ and $\lambda = 810\text{nm}$, yielding $\omega_\perp \simeq 2\pi \times 38\text{kHz}$. For $N_a = 1.4 \times 10^5$ atoms [13], $R_x \simeq 14\mu\text{m}$, $R_y \simeq 20\mu\text{m}$, $R_z \simeq 10.6\mu\text{m}$, and within the classical radius there are $\pi R_x R_y / d^2 \simeq 5400$ atom tubes. From Eq. (1) we obtain the number of atoms in each tube $N(x_0, y_0) = \int d^3r \rho = 5N_0 d^2 (1 - x_0^2/R_x^2 - y_0^2/R_y^2)^{3/2} / 2\pi R_x R_y$. At the central tube, $N(0, 0) \simeq 65$ atoms.

When a shallow optical lattice is applied, the wave function Eq. (1) is modified along the z axis. In the experiment the displacement of the harmonic potential was about $\delta \simeq 3\mu\text{m} \simeq 2.2l$, where $l \equiv (\hbar/m\omega)^{1/2}$ and the trap frequency $\omega \simeq 2\pi \times 60\text{Hz}$. Within $2R_z$, there are about 52 lattice sites.

We study the NIST experiment of dipolar oscillations within the TWA, by suddenly displacing the harmonic trap. In the TWA, in a tight elongated trap $\omega \ll \omega_\perp$, the dynamics of the 1D classical stochastic field $\psi_W(z, t)$ follows from the nonlinear equation [19]:

$$i \frac{\partial}{\partial t} \psi_W = \left(-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial z^2} + V + g|\psi_W|^2 \right) \psi_W, \quad (2)$$

which coincides with the Gross-Pitaevskii equation (GPE). Here the potential is a combined harmonic trap and a periodic optical lattice $V(x) = m\omega^2 z^2/2 + sE_r \sin^2(\pi z/d)$ and $g = 2\hbar\omega_\perp a$, where a denotes the scattering length. The thermal and quantum fluctuations are included in the initial state of ψ_W in Eq. (2) which represents an ensemble of Wigner distributed wave functions. Our basic formalism of the TWA is similar to Ref. [15]. Initially, before displacing the trap, the gas is assumed to be in thermal equilibrium and we approximate the field operator $\hat{\psi}(z, t=0)$, within the Bogoliubov theory [20]:

$$\hat{\psi}(z) = \psi_0(z)\hat{\alpha}_0 + \sum_{j>0} [u_j(z)\hat{\alpha}_j - v_j^*(z)\hat{\alpha}_j^\dagger]. \quad (3)$$

Here ψ_0 is the ground state solution of the GPE, obtained by evolving the GPE in imaginary time, and $\langle \hat{\alpha}_0^\dagger \hat{\alpha}_0 \rangle = N_0$, the number of ground state atoms. The mode functions $u_j(z)$ and $v_j(z)$ ($j > 0$), and the energies ϵ_j for excited states are obtained by solving the corresponding Bogoliubov equations [15] in the harmonic trap, modulated by the shallow lattice potential. In the TWA the time evolution of the ensemble of Wigner distributed wave functions [Eq. (2)] is unraveled into stochastic trajectories, where the initial state of each realization for ψ_W is generated according to Eq. (3) with the quantum operators $(\hat{\alpha}_j, \hat{\alpha}_j^\dagger)$ replaced by complex Gaussian-distributed random variables (α_j, α_j^*) . These are obtained by sampling the corresponding Wigner distributions [15]. In particular, for the quasiparticles the

Wigner function is that of the ideal harmonic oscillators in a thermal bath [21], with the width $\bar{n}_j + \frac{1}{2}$, where $\bar{n}_j \equiv \langle \hat{\alpha}_j^\dagger \hat{\alpha}_j \rangle = [\exp(\epsilon_j/k_B T) - 1]^{-1}$ and the $1/2$ width at $T = 0$ for each mode represents the quantum noise.

Because the Wigner function returns symmetrically ordered expectation values, $\langle \alpha_j^* \alpha_j \rangle_W = \bar{n}_j + \frac{1}{2}$ and $\langle \alpha_j \rangle_W = \langle \alpha_j^* \rangle_W = \langle \alpha_j^2 \rangle_W = 0$, for $j > 0$, it is generally not possible to extract from the TWA simulations the correlation functions for the full multi-mode field operator [19]. Consequently, we derive the desired normally ordered expectation values by defining the ground state operators a_j for each individual lattice site j [15]: $a_j(t) \equiv \int dz \psi_0^*(z) \psi_W(z, t)$, where the integration is over the j th lattice site, $\psi_W(z, t)$ is the stochastic field, determined by Eq. (2), and $\psi_0(z)$ is the ground state wave function, obtained by evolving the GPE in imaginary time. This provides us with the basis for $\tilde{\psi}(z) = \sum_j \hat{a}_j \phi_j(z)$, where the ground state functions ϕ_j are restricted in the j th site, and the normally ordered expectation values are easily obtained with respect to $\tilde{\psi}$, e.g., for the cm position z_{cm} , the position fluctuations $\Delta z = [\langle \hat{z}^2 \rangle - \langle \hat{z} \rangle^2]^{1/2}$, and for the normalized phase coherence between the central well and its i th neighbor $C_i \equiv |\langle \hat{a}_0^\dagger \hat{a}_i \rangle| / \sqrt{n_0 n_i}$.

We numerically study the dipole oscillations using the experimental parameters [13]. In the central atom tube $N \simeq 65$, resulting in the nonlinearity $Ng \simeq 320\hbar\omega l$ for $a \simeq 5.313\text{nm}$. In Fig. 1 we use the same fixed Ng , but vary N , and show the cm position z_{cm} of the atoms for $s = 0.25$. The atoms are initially in thermal equilibrium at $t = 0$ and at $\omega t_0 = 1$ we displace the center of the harmonic trap from $z = 0$ to $z = \delta = 2.16l \simeq 7.4d$. In the initial state (3), we generate the noise for 300 modes, corresponding to 4-5 lowest energy bands. Even though the atoms mainly remain in the lowest band, synthesizing the TWA noise for only 70 modes ($\sim$ one band) can underestimate the damping and the phase decoherence several tens of percents. This emphasizes the importance of the multi-mode approach to the TWA, beyond the tight-binding approximation, and is consistent with our previous observations [15]. In 1D TWA, unlike in 3D [22], we did not find any divergence of the calculated results, when the mode number was increased. For each run, we typically use 600 stochastic realizations. The integration of the nonlinear evolution in Eq. (2) is obtained using the split-step method [23] on a spatial grid of 4096 points. The numerics becomes much faster for larger atom numbers, due to smaller quantum fluctuations.

As in Ref. [13], we model the cm motion as a damped harmonic oscillator $\ddot{z}_{\text{cm}} = -kz/m^* - 2\gamma\dot{z}$, so that for underdamped motion $z_{\text{cm}}(t') \equiv -e^{-\gamma t'}(\delta \cos \Omega t' + c \sin \Omega t')$, with $c \equiv \gamma\delta/\Omega$ and $\Omega = \sqrt{k/m^* - \gamma^2}$, where $t' \equiv t - t_0$ and m^* denotes the effective mass. At $t' = 0$ we have $z_{\text{cm}} = -\delta$ with $v_{\text{cm}} = 0$. Even in a very shallow lattice, $s = 0.25$, we find a significant damping for $N = 75$ and 90 atoms, with $\gamma/\omega \simeq 0.020$ and 0.013 [Fig. 1]. The damp-




FIG. 1: The oscillations of the cm position of atoms from the harmonic trap center at $T = 0$ with fixed Ng (top left; curves from top $N = 400, 125, 75$), the damping rates (top right) $\gamma(N)$ and $\gamma_v(N)$ (upper line), and the normalized quasimomentum distributions at $\omega t = 8$ (bottom left) with shaded unstable region; the inset shows the population $n(p)\hbar/l$ at $v = 9\omega l$ where the classical motion becomes unstable. The trap is instantaneously displaced at $\omega t = 1$. The phase coherence between the initially central well and its fifth neighbor C_5 (bottom right; curves from top $N = 1000, 400, 125, 90, 75$).

ing is clearly correlated to the loss of phase coherence. The dependence of γ on N is approximately exponential $\gamma(N) \simeq 0.11\omega \exp(-N/43)$, resulting in $\gamma \simeq 0.025\omega$ at the central tube with $N \simeq 65$, which is close to the experimentally observed value of $\gamma \simeq 0.03\text{--}0.04(\pm 0.01)\omega$ [13]. We also checked that the damping rate γ_v , obtained from $v_{\text{cm}} = dz_{\text{cm}}/dt$, is close to γ . For $s = 0.25$ the single-particle effective mass is almost equal to m and we find $\Omega \simeq \omega$, with very little effects of fluctuations in Ω .

In the experiment [13] the dynamics was measured by imaging the expanding atom cloud, indicating that γ did not represent the value in the central tube, but the average value over all the tubes. Although in the experiment g is constant in each tube (as N varies), we can obtain an upper bound limit for the average damping rate by extrapolating our fitted $\gamma(N)$ (for fixed Ng) to small N , so that for $\gamma/\omega \ll 1$, $\bar{\gamma} = \sum_{i,j} \gamma(N)N(x_i, x_j)/N_a$. Here $N(x_i, x_j)$ is the atom number in a tube at (x_i, y_j) and the summation is over all the tubes. By replacing the sum by an integral, we obtain $\bar{\gamma} \simeq 0.047\omega$.

We also studied thermal effects on the dipolar motion [Fig. 2]. In the experiments [13] it was not possible to derive information about temperature and the effect of thermal atoms on γ . For small N we found thermal fluctuations to be negligible compared to quantum fluctuations. Only for ground state atom numbers clearly exceeding 400, we can recognize any notable finite-T effects on γ . This strongly suggests that the experimentally observed damping [13] duly was due to quantum fluctuations.

The classical nonlinear evolution of a BEC in a periodic lattice is susceptible to dynamical instabilities arising from both constant velocity, larger than a critical value [3, 4, 5, 6], and acceleration or force, smaller than




FIG. 2: Thermal effects on the coherence and the cm motion. The coherence C_5 , as in Fig. 1 (left); curves from top represent $(N_0, k_B T/\hbar\omega) = (1000, 0)$, $(900, 21.3)$, $(400, 0)$, and $(360, 10.5)$. The cm (right) for the same cases. Only the finite-T, $N_0 = 900$ case has notable damping ($\gamma \simeq 0.002\omega$).

a critical value [7, 8, 9]. The onset of the instability is associated with inhibited transport, fragmentation of the density profile, and the significant broadening of quasimomentum distribution. In Ref. [13] the importance of the dynamical instabilities on dissipation was not clear, since the velocity was much smaller than the critical value and no momentum broadening was observed, despite the strongly damped motion.

In the combined harmonic and lattice potential, the separation of the acceleration and the velocity instabilities is not straightforward, since the trap induces a local force $m\omega^2 z$ and, e.g., applying a constant force $m\omega^2 F$ on atoms is the same as displacing the trap: $z^2/2 - zF = (z - F)^2/2 + \text{const}$. Moreover, a weakly accelerating lattice does not equal to applying a constant force, since the acceleration in a shallow lattice ($s = 0.25$) does not induce notable cm motion. We numerically integrated the classical GPE for a BEC (without quantum or thermal noise) and found the onset of the dynamical instability at $\delta_c \simeq 8\text{--}9l$, corresponding to a maximum velocity $v_{\text{cm}}^c \simeq 8\text{--}9\omega l$. For smaller displacements, no instabilities were observed on the time scale of the experiment [13], e.g., due to the weak force induced by the trap. With quantum fluctuations, the classically observed sharp onset of the dynamical instability is smeared out due to the phase uncertainty of atoms between neighboring lattice sites [14]. This is similar to a BEC in a double-well potential, where the macroscopic self-trapping of the atom population can decay due to dissipation [24].

In order to investigate the role of the classical dynamical instabilities in the dissipative quantum dynamics, we calculated the quasimomentum distribution in the TWA from $\hat{\psi}$ and found atoms occupying the dynamically unstable velocity region of the corresponding classical system; Fig. 1. This population is larger and grows more rapidly, the larger the observed damping: e.g., at $v = 9\omega l$ and $\omega t = 8$, $n(p)\hbar/l \simeq 0.02$ and 0.08 , for $N = 225$ and 75 , and both $n(p)$ and γ show similar exponential dependence on N . For $(N, Ng/\hbar\omega l) = (90, 450)$ (here $\gamma \simeq 0.026\omega$), the ratio of the population at the unstable velocity $v \simeq 9\omega l$ to that at the peak of the distribution was 0.07 at $\omega t = 3$ and as high as 0.09 at $\omega t = 6$. It is this finite high velocity occupation that results in dissi-




FIG. 3: The coherence C_5 (left) in a displaced trap, in a lattice moving with a constant velocity $2.16l\omega$, in an accelerated ($0.27l\omega^2$ and $0.54l\omega^2$) lattice, and in a static lattice. Only the $0.54l\omega^2$ accelerated lattice exhibits a lower C_5 . The quasimomentum distribution (right) in a displaced trap (off-centered curve) and in a static potential at $\omega t = 6$.

pation. Alternatively, we may say that the position uncertainty Δz due to vacuum fluctuations is comparable to the critical displacement δ_c , resulting in dissipation, even though $\delta < \delta_c$. In the example case $\Delta z \simeq 5-6l$. This is consistent with the observation in Ref. [16] that the damping arises from the large depletion, due to the tight transverse confinement, which leads to the population of high-momentum states. The tails of the momentum distribution can become very long when the atoms reach the 1D Tonks-Girardeau regime [25].

Moreover, we also studied the atom response to an accelerated lattice, to a lattice moving with a constant velocity, and to a static lattice, without displacing the trap; see Fig. 3. We used the parameters of Fig. 1, with $N = 125$ and $Ng = 405$. The constant velocity was $v = 2.16l\omega$ and the acceleration $0.27l\omega^2$ over the time period of $\omega t = 8$, so that the final velocity was $2.16l\omega$. In all the cases z_{cm} remained close to zero, without notable cm motion. The decrease in C_i in all the three systems was almost the same as for the displaced trap, indicating that the motion of neither the atoms, nor the lattice, had significant effect on the phase coherence in the experiments. The occupation of the dynamically unstable region in the quasimomentum distribution in the static potential was approximately equal to that in the displaced trap case (e.g., at $p = \pm 9\hbar/l$, $n \simeq 0.05l/\hbar$), resulting in a similar number of high velocity atoms in the two cases. Moreover, the variation of this atom number was very little affected by the dipole oscillations. Consequently, neither phase decoherence nor the population with unstable velocities, which result in the damping of the dipole oscillations, are here due to the motion of the atoms, but the combined effect of the strong transverse confinement and the lattice. This is contrary to the system studied in Ref. [14] where the atom motion itself was argued to generate the decoherence.

As expected, both the increase of the lattice height and the initial displacement enhance dissipation. We studied the dynamics for $s = 1$, $\delta = 2.16l$, and obtained for $(N, Ng/\hbar\omega l) = (330, 350), (430, 340), (830, 330)$, $\gamma/\omega \simeq 0.019, 0.012, 0.003$, at $T = 0$. Also with $s = 1$ the damping is notable for δ much smaller than the classical critical

displacement $\delta_c \simeq 6-7l$. On the other hand, for $N = 175$ and $Ng \simeq 380\hbar\omega l$, the increase of δ/l from 2.16 to 4.32 increased γ/ω from 0.004 to 0.009.

In summary, we studied dissipative atom transport in a shallow 1D lattice. At a fixed nonlinearity, smaller N results in larger phase and momentum uncertainty and, consequently, larger population in the dynamically unstable velocity region. Surprisingly, this population is generated due to the confinement and the lattice, even with neither the atoms nor the lattice moving. The qualitative agreement between the TWA and the experiment [13], despite the approximations involved in generating the initial state at small N , also represents a considerable success of the TWA which is traditionally considered more suitable in the limit of large field amplitude [19, 22]. For instance, the time-dependent Hartree-Fock-Bogoliubov theory predicts much too small a damping [16]. This suggests the TWA could provide a powerful technique to study vacuum fluctuations also in several other related condensed matter and atomic systems, such as corrugated superconducting nanowires [26].

We acknowledge financial support from the EPSRC and NSF through a grant for the Institute for Theoretical Atomic, Molecular and Optical Physics at Harvard University and Smithsonian Astrophysical Observatory. After our work was completed, the dynamics in a deep lattice was analyzed in the fermionized limit [27], in the Mott state. In this limit the TWA with a bosonic field amplitude is no longer valid.

-
- [1] S. Burger *et al.*, Phys. Rev. Lett. **86**, 4447 (2001).
 - [2] F. Ferlaino *et al.*, Phys. Rev. A **66**, 011604 (2002).
 - [3] B. Wu and Q. Niu, Phys. Rev. A **64**, 061603 (2001).
 - [4] A. Smerzi *et al.*, Phys. Rev. Lett. **89**, 170402 (2002).
 - [5] F.S. Cataliotti *et al.*, New J. Phys. **5**, 71 (2003).
 - [6] L. Fallani *et al.*, Phys. Rev. Lett. **93**, 140406 (2004).
 - [7] M. Cristiani *et al.*, Opt. Express **12**, 4 (2004).
 - [8] Y. Zheng *et al.*, Phys. Rev. Lett. **93**, 230401 (2004).
 - [9] A.R. Kolovsky, e-print cond-mat/0412195.
 - [10] L. De Sarlo *et al.*, e-print cond-mat/0412279.
 - [11] M. Greiner *et al.*, Nature **415**, 39 (2002); T. Stöferle *et al.*, Phys. Rev. Lett. **92**, 130403 (2004); B. Paredes *et al.*, Nature **429**, 277 (2004).
 - [12] D. Jaksch *et al.*, Phys. Rev. Lett. **81**, 3108 (1998).
 - [13] C.D. Fertig *et al.*, Phys. Rev. Lett. **94**, 220402 (2005).
 - [14] A. Polkovnikov and D.-W. Wang, Phys. Rev. Lett. **93**, 070401 (2004).
 - [15] L. Isella and J. Ruostekoski, e-print cond-mat/0409475.
 - [16] J. Gea-Banacloche *et al.*, e-print cond-mat/0410677.
 - [17] E. Altman *et al.*, cond-mat/0411047.
 - [18] B.L. Tolra *et al.*, Phys. Rev. Lett. **92**, 190401 (2004).
 - [19] M.J. Steel *et al.*, Phys. Rev. A **58**, 4824 (1998).
 - [20] 1D Bogoliubov becomes progressively worse at small N , due to increasing density fluctuations: K.V. Kheruntsyan *et al.*, Phys. Rev. Lett. **91**, 040403 (2003).
 - [21] C.W. Gardiner and P. Zoller, *Quantum Noise* (Springer,

- Berlin, 1999).
- [22] A. Sinatra *et al.*, J. Phys. B **35**, 3599 (2002).
 - [23] J. Javanainen and J. Ruostekoski, cond-mat/0411154.
 - [24] J. Ruostekoski and D.F. Walls, Phys. Rev. A **58**, R50 (1998).
 - [25] M. Olshanii, Phys. Rev. Lett. **81**, 938 (1998).
 - [26] A. Bezryadin *et al.*, Nature **404**, 971 (2000).
 - [27] A.M. Rey *et al.*, cond-mat/0503477; M. Rigol *et al.*, cond-mat/0503302; G. Pupillo *et al.*, cond-mat/0505325.